



EAAP + WAAP + Interbull Congress 2023
 Lyon, France - August 26th / September 1st, 2023

Monitoring liveweight in Sarda dairy sheep using a walk-over-weighing system

M. Decandia¹, M. Acciaro¹, V. Giovanetti¹, G. Molle¹, F. Chessa¹, I. Llach², E. González-García²
¹AGRIS Sardegna, Loc. Bonassai S.S. 291 Sassari-Fertilia - Km. 18,600 – Sassari (Italy)
²SELMET, INRAE, Montpellier SupAgro, CIRAD, Université Montpellier, (France)

Liveweight (**LW**) monitoring is important in livestock management to check nutritional, reproduction and welfare status of animals. In dairy sheep, LW measurement is less frequent because is time and labor consuming. The use of automated weighing system could make easier this practice.

Within the H2020 TechCare project, aiming to test innovative technologies to improve welfare management for small ruminant systems, a short-test was run to evaluate the precision and accuracy of LW measures carried out using an automated walk-over-weighing (**WoW**) scale (González-García *et al.*, 2018), which was compared with a static scale.

Thirty-six dry Sarda dairy ewes (LW, mean±SE 55.49 ±0.92) were used in a 3-day session. Ewes were daily weighed 3 times following the same circuit, which included a first static LW measurement; once a static position was achieved and LW recorded, the animals continued and traversed the WoW scale for the WoW LW recording. The ewes were previously accustomed to the circuit during a pre-experimental week (see Figure below).



Raw static and automatic LW data were first filtered for removing misbehaviors and outlier with Kalman filter algorithm (Cloeze *et al.*, 2022), using the ORIOLE web application (<https://oriole.sk8.inrae.fr/>)

The concordance correlation coefficient (CCC; Lin, 1989) has been then calculated (Model Evaluation System; Tedeschi, 2006) to simultaneously account for precision and accuracy. The two component of CCC (=0.98), correlation coefficient estimate (*r*, that measure precision) and the bias correction factor (*C_b*, that indicates the accuracy) were respectively 0.98 and 0.99 indicating high agreement between measures.

The WoW system evaluated here is an alternative to the static scales conventionally used on dairy sheep farms. If sound filtration of raw data is applied, WoW could contribute to the close (daily) monitoring of individual LW without operator intervention (i.e., voluntary weighing), taking animal welfare into account (i.e., no stress related to the weighing session on static scales), and potentially detecting nutrition issues highlighted by LW changes.



@TechCareproject



@TechCareproject



@TechCareproject

The EU support of Techcare project is acknowledged

www.techcare-project.eu



Integrating innovative TECHNOLOGIES along the value Chain to improve small ruminant welfare management